

Piecewise linear loop quantum gravity

Jonathan Engle

Albert-Einstein-Institut, Am Mühlenberg 1, D-14476 Golm, EU

December 6, 2008

Abstract

We define a modification of LQG in which graphs are required to consist in piecewise linear edges, which we call piecewise linear LQG (pLQG). At the diffeomorphism invariant level, we prove that pLQG is equivalent to standard LQG, as long as one chooses the class of diffeomorphisms appropriately. That is, we exhibit a unitary map between the diffeomorphism invariant Hilbert spaces that maps physically equivalent operators into each other. In addition, using the same ideas as in standard LQG, one can define a Hamiltonian and Master constraint in pLQG, and the unitary map between pLQG and LQG then provides an exact isomorphism of dynamics in the two frameworks.

Furthermore, loop quantum cosmology (LQC) can be exactly embedded into pLQG. This allows a prior program of the author to embed LQC into LQG at the dynamical level to proceed. In particular, this allows a formal expression for a physically motivated embedding of LQC into LQG at the diffeomorphism invariant level to be given.

1 Introduction

Loop quantum gravity (LQG) [1, 2, 3] is a minimalistic, background independent approach to quantum gravity. However, in the construction of the theory, technical choices have to be made, especially in the kinematics of the theory. One can then ask: might some of these technical choices not matter once the constraints are solved? In this paper, we show that in particular the choice of the piecewise analytic category is not essential: it can even be replaced with something as simple as the piecewise linear category, *and the resulting theory is the same at the diffeomorphism invariant level*. The diffeomorphism invariant Hilbert spaces of the two theories are naturally isomorphic, and the dynamics are exactly the same. Furthermore, a very large algebra of the diffeomorphism invariant operators are also seen to be the same.

We call this modification of LQG ‘piecewise linear loop quantum gravity’ (pLQG).

What are the consequences of this? First, this can be used as a “trick” to circumvent the obstruction to the program of [4, 5] caused by the non-embeddability result proved in [6]. One can arrive at the same diffeomorphism invariant Hilbert space as in standard LQG, but by means of a piecewise linear kinematics that completely circumvents [6]. As a result, it is possible to write down formal expressions for embeddings of LQC into LQG at the diffeomorphism invariant level, of the type systematically motivated in the work [4, 5]. Second, this new framework might allow a closer relation to spinfoams [8, 9], which also use the piecewise linear category to define the kinematics [9].

After this work was completed, it was pointed out to the author that the kinematics of piecewise linear LQG as presented here, and the choice of generalized diffeomorphisms, had already been proposed as a model in [7]. However [7] was not interested in pLQG as such, and so did not develop it beyond kinematics. This paper goes further, in rigorously constructing the rigging map for the

diffeomorphism constraint, constructing Hamiltonian and Master constraint operators, and showing equivalence with the piecewise analytic framework at the diffeomorphism invariant level including dynamics. Of course the embedding of LQC into pLQG is also new. On the other hand, [7] presents features of the kinematics of pLQG not presented here. For example, [7] introduces the piecewise linear analogue $\overline{\mathcal{A}}_{PL}$ of the generalized connections, and constructs the piecewise linear analogue of the Ashtekar-Lewandowski measure, allowing one to express the kinematical Hilbert space as an L^2 space.

The paper is organized as follows. First we define the kinematics of piecewise linear LQG, motivate a choice of generalized diffeomorphism group, and solve the diffeomorphism constraint. The unitary map between the diffeomorphism invariant Hilbert spaces in pLQG and LQG is then explicitly constructed and proven in section 3. Equivalence of diffeomorphism invariant operators in the two frameworks, and the equivalence of dynamics in the two frameworks is proven in section 4. The exact embeddings of LQC into pLQG of the type motivated in [4, 5] are then explicitly reviewed in section 5, and at the end of this section, the resulting formal expressions for the embeddings of LQC into LQG at the diffeomorphism invariant level are given. We then close with a brief discussion.

2 Piecewise linear loop quantum gravity

2.1 Kinematics

We assume space, M , is topologically $\mathbb{R}^3$, and we equip M with a fixed, flat frame bundle connection ∂_a . This flat connection gives us a notion of ‘straightness’ on M .

Let $\mathcal{A}$ denote the space of smooth $SU(2)$ connections on M . The classical phase space is parametrized by such a connection A_a^i and a densitized triad field $\tilde{E}_i^a$. (Here A_a^i denotes the components of the $SU(2)$ connection with respect to the basis $\tau_i := -\frac{i}{2}\sigma_i$ of the Lie algebra $\mathfrak{su}(2)$.) The Poisson brackets are given by

$$\{A_a^i(x), \tilde{E}_j^b(y)\} = 8\pi\gamma G \delta_j^i \delta_a^b \delta^3(x, y) \quad (1)$$

where G is Newton’s constant, and $\gamma \in \mathbb{R}^+$ is the Barbero-Immirzi parameter.

Next one specifies the basic variables. The algebra of elementary configuration variables is chosen to consist in (real analytic¹) functions of finite numbers of holonomies of the connection A_a^i along piecewise *straight* edges; we will also use the term *piecewise linear* for such edges. We call these functions *piecewise linear cylindrical* and the space of such functions is denoted $\underline{\text{Cyl}}$. The elementary momentum variables are taken to be the *fluxes* on piecewise flat surfaces². Given a surface S and a function $f : S \rightarrow \mathfrak{su}(2)$, we define the corresponding flux by

$$E(S, f) := \int_S f^i \tilde{E}_i^a n_a d\sigma_1 d\sigma_2 \quad (2)$$

where $n_a := \epsilon_{abc} \frac{\partial x^b}{\partial \sigma_1} \frac{\partial x^c}{\partial \sigma_2}$, (σ_1, σ_2) are arbitrary coordinates on S , x^a are arbitrary coordinates on the spatial manifold, and ϵ_{abc} denotes the fully anti-symmetric symbol (i.e., the Levi-Civita tensor of density weight -1).

¹ As always, one has some freedom in the precise definition of cylindrical functions. This is the definition that will be convenient for this paper.

²We may also include the fluxes on arbitrary piecewise analytic surfaces, but nothing is thereby gained, and using piecewise flat surfaces is more in the spirit of piecewise linear loop quantum gravity as presented here.

Next let us introduce some structures to give a more useful characterization of $\underline{\text{Cyl}}$. We first define a *piecewise linear path* to be a continuous path $e : [0, 1] \rightarrow M$ consisting in a finite number of segments, each segment being geodesic with respect to ∂_a (but not necessarily affinely parametrized.) We then define a *piecewise linear edge* to be an equivalence class of piecewise linear paths, where two piecewise linear paths are equivalent if they are related by a reparametrization, or addition or removal of ‘trivial’ segment of the form $(\delta \circ \delta^{-1})$.³ We next define a *piecewise linear graph* to be a finite, ordered set of piecewise linear edges. Let $\underline{\Gamma}$ denote the space of piecewise linear graphs. With these definitions, any element Φ of $\underline{\text{Cyl}}$ can be written in the form

$$\Phi[A] = F(A(e_1), \dots, A(e_n)) \quad (3)$$

for some piecewise linear graph $(e_1, \dots, e_n) \in \underline{\Gamma}$, and some real-analytic function $F : SU(2)^n \rightarrow \mathbb{C}$. If a cylindrical function $\Phi \in \underline{\text{Cyl}}$ may be written using the edges of a graph γ , we say Φ is *cylindrical with respect to γ* . We denote by Cyl_γ the space of functions cylindrical with respect to γ .

We next define an inner product $\langle \cdot, \cdot \rangle$ on $\underline{\text{Cyl}}$ in the same way as in standard LQG: Given $\Psi, \Phi \in \underline{\text{Cyl}}$, we find a graph γ large enough so that $\Psi, \Phi \in \text{Cyl}_\gamma$, and then define the inner product between Ψ and Φ using the Haar measure on $SU(2)$. As in LQG, this inner product is independent of the ambiguity in the choice of γ . For each γ let $\mathcal{H}_\gamma$ denote the Cauchy completion of Cyl_γ , and let $\underline{\mathcal{H}}$ denote the Cauchy completion of $\underline{\text{Cyl}}$, in this inner product.

We next construct a representation of the basic algebra on $(\underline{\text{Cyl}}, \langle \cdot, \cdot \rangle)$. The configuration algebra $\underline{\text{Cyl}}$ is represented by multiplication. The operators corresponding to the momentum degrees of freedom are then defined via the classical Poisson bracket

$$\widehat{E(S, f)}\Phi = i\{E(S, f), \Phi\} \quad (4)$$

which ensures that the commutators of elements of $\underline{\text{Cyl}}$ and the fluxes match the corresponding Poisson brackets correctly. The multiplicative $\underline{\text{Cyl}}$ operators are bounded because each element of $\underline{\text{Cyl}}$, as a continuous function of a finite number of $SU(2)$ holonomies, is bounded due to the compactness of $SU(2)$. These multiplicative operators thus extend to all of $\underline{\mathcal{H}}$ by the BLT theorem. The flux operators, equipped with domain $\underline{\text{Cyl}}$, form essentially self-adjoint operators, which therefore extend uniquely to self-adjoint operators on $\underline{\mathcal{H}}$. One can check that the resulting representation of the basic observables then reflects correctly not only the poisson brackets, but also the correct adjointness relations. This is the elementary quantization.

After the quantization of the elementary operators, other geometrical operators corresponding to length, area, and volume can also be constructed in the same way as in standard LQG [1, 2, 10], all with the same spectra. The Gauss constraint is defined in the same way as in standard LQG [1, 2] and is just as easy to solve, yielding as a solution space $\underline{\mathcal{H}}_G \subset \underline{\mathcal{H}}$, consisting in the Cauchy completion of the span of gauge-invariant spin-network states [1, 2], but this time restricted to graphs in $\underline{\Gamma}$.

2.2 Solution to the diffeomorphism constraint

Next, let us discuss the solution to the diffeomorphism constraint. Central to this is the selection of a generalization of the group of diffeomorphisms to be used in quantum theory. Once this generalization is selected, we will simply use the group averaging strategy of [1, 11] to solve the constraint.

The choice of diffeomorphism gauge group

³Thus, two paths are equivalent iff they allows yield the same holonomies.

Let $\underline{\text{Diff}}$ denote the group of generalized diffeomorphisms to be used. We first stipulate several requirements of $\underline{\text{Diff}}$, which will lead us to a choice for the group. First, we stipulate that the generalized diffeomorphisms at least consist in bijective maps of space onto itself.⁴ Second, each element of $\underline{\text{Diff}}$ must map all piecewise linear edges to piecewise linear edges, so that it has a well-defined action on $\underline{\Gamma}$, the set of piecewise linear graphs. These requirements, however, so far are not enough: if we were to only require these, one could map any graph into any other with such a ‘generalized diffeomorphism’, and, if one follows the prescription of [1, 11], one would be led to a solution space with only a single state. Therefore, we furthermore stipulate that the maps be homeomorphisms. A natural choice satisfying the above requirements is the group of *piecewise linear homeomorphisms*. To define the notion of a piecewise linear homeomorphism, we must first review the definition of a simplicial complex [12]. First, we note that the fixed connection ∂_a endows M with a natural *affine structure*. Let us for convenience arbitrarily pick an origin $O \in M$, and use this to make M into a vector space, so that addition and real scalar multiplication are defined in M . None of the definitions or constructions below will depend on the choice of O .

A set of points $\{a_0, \dots, a_n\} \subset M$ is said to be *independent* if they do not lie within any common $(n-1)$ -dimensional plane in M . Given such a set of $n+1$ independent points, we define the *n-simplex* σ spanned by $a_0, \dots, a_n$ to be the set of all points $x \in M$ such that

$$x = \sum_{i=0}^n t_i a_i \quad (5)$$

for some $t_0, \dots, t_n \in \mathbb{R}$ all non-negative, satisfying $\sum_{i=0}^n t_i = 1$. n is called the *dimension* of σ . In common language, a 0-simplex is a point, a 1-simplex is a line segment, a 2-simplex is a triangle, and a 3-simplex is a tetrahedron.

Next we define the generalized notion of ‘face’. Given an n -simplex σ spanned by a set of points $\{a_0, \dots, a_n\}$, the simplex spanned by a subset of these points is called a *face* of σ . In particular, every simplex is a face of itself; a face of a simplex σ that is not equal to σ is called a *proper face*. Thus, in this generalized sense, the proper ‘faces’ of a tetrahedron consist in all the triangular faces in the usual sense, all the edges, and all four vertices. The proper ‘faces’ of a triangle consist in its three edges and three vertices, etc.

We can now define a *simplicial complex* K to be a (possibly infinite) collection of simplices such that

1. Every face of a simplex of K is in K .
2. The intersection of any two simplices of K is a face of each of them.

The maximal simplex dimension occurring in K is called the *dimension of K* .

Finally, a homeomorphism F from an n -dimensional manifold M onto an n -dimensional manifold N is called a *piecewise linear* if there exist simplicial complexes K and L , covering all of M and N , respectively, such that $v_1, \dots, v_m$ span a simplex of K if and only if $F(v_1), \dots, F(v_m)$ span a simplex of L , and such that for each $\{v_0, \dots, v_n\}$ spanning an n -simplex in K ,

$$F\left(\sum_{i=0}^n t_i v_i\right) = \sum_{i=0}^n t_i F(v_i) \quad (6)$$

⁴If one were to solve the Gauss and diffeomorphism constraints together, this would be equivalent to requiring that the generalized principal bundle automorphisms to be used should at least consist in maps from the principal bundle to itself that preserve all structure of the principal bundle except possibly topology and differentiable structure.

for all $t_i \geq 0$ satisfying $\sum_{i=0}^n t_i = 1$. Said simply, F maps simplices of K into simplices of L in a continuous way, such that F is linear within each n -simplex.⁵

The piecewise linear homeomorphisms are essentially the piecewise linear analogue of the *stratified analytic diffeomorphisms* advocated in [13] and described in [14, 15] (see also [16]). In the analytic framework, however, one has more choices: one can, for example, require that the generalized diffeomorphisms be at least differentiable. The analogue of such a requirement can, however, *not* be satisfied in the piecewise linear framework: the only differentiable piecewise linear maps are fully linear. But the group of globally linear maps is too small: if one were to choose $\underline{\text{Diff}}$ to be the group of linear maps, even individual open edges would have global information that would be diffeomorphism-invariant. This would prevent any possible relation, with any analytic LQG framework so far proposed, at the diffeomorphism-invariant level.

Construction of the diffeomorphism invariant Hilbert space

With the foregoing choice of $\underline{\text{Diff}}$, let us proceed to construct the solution to the diffeomorphism constraint. For this purpose, we introduce some further definitions. First, if two graphs $\gamma_1, \gamma_2 \in \underline{\Gamma}$ differ only by a permutation of edges or reversal of edge orientations, call them *probe equivalent*. The probe equivalence class of a graph γ we write as $[\gamma]_{\text{pr}}$. Let $\underline{\Gamma}_{\text{pr}}$ denote the space of such probe equivalence classes in $\underline{\Gamma}$. Next, for each $\gamma \in \underline{\Gamma}$, let $\mathcal{H}'_\gamma$ denote the orthogonal complement, in $\mathcal{H}_\gamma$, of the span of all functions that are constant on at least one edge of γ . Then, as in [1],

$$\underline{\mathcal{H}} = \oplus_{[\gamma]_{\text{pr}} \in \underline{\Gamma}_{\text{pr}}} \mathcal{H}'_\gamma. \quad (7)$$

Furthermore let $\text{Cyl}'_\gamma := \mathcal{H}'_\gamma \cap \text{Cyl}$. Lastly we define some subgroups of our chosen generalized diffeomorphisms. For each $\gamma \in \underline{\Gamma}$, let $\underline{\text{Diff}}_\gamma$ be the set of elements in $\underline{\text{Diff}}$ mapping γ back into its probe equivalence class. Let $\underline{\text{TDiff}}_\gamma$ be the set of elements in $\underline{\text{Diff}}$ fixing γ , so that they preserve each edge of γ including orientation. Let $\underline{GS}_\gamma := \underline{\text{Diff}}_\gamma / \underline{\text{TDiff}}_\gamma$ where the division is taken with respect to the left-action.

For each $\gamma \in \underline{\Gamma}$, define $\underline{P}_{\text{diff}, \gamma}$ as the *group averaging map* [11, 1] from $\mathcal{H}'_\gamma$ to the subspace invariant under $\underline{GS}_\gamma$.⁶

$$\underline{P}_{\text{diff}, \gamma} \Psi_\gamma := \frac{1}{|\underline{GS}_\gamma|} \sum_{\varphi \in \underline{GS}_\gamma} \varphi^* \Psi_\gamma. \quad (8)$$

For each $\Psi_\gamma \in \text{Cyl}'_\gamma$, define $\underline{\eta}(\Psi_\gamma) \in \underline{\text{Cyl}}^*$ by

$$(\underline{\eta}(\Psi_\gamma)|\Phi) := \sum_{\varphi \in \underline{\text{Diff}}/\underline{\text{Diff}}_\gamma} \langle \varphi^* \underline{P}_{\text{diff}, \gamma} \Psi_\gamma, \Phi \rangle = \frac{1}{|\underline{GS}_\gamma|} \sum_{\varphi \in \underline{\text{Diff}}/\underline{\text{TDiff}}_\gamma} \langle \varphi^* \Psi_\gamma, \Phi \rangle. \quad (9)$$

Piecing together these maps for the various $\gamma \in \underline{\Gamma}$ defines a map $\underline{\eta} : \underline{\text{Cyl}} \rightarrow \underline{\text{Cyl}}^*$. This is the rigging map for solving the diffeomorphism constraint for piecewise linear LQG. The space of ‘test functions’ at the diffeomorphism invariant level is then

$$\underline{\text{Cyl}}^*_{\text{diff}} := \text{Im} \underline{\eta}. \quad (10)$$

⁵In the language of [12], a piecewise linear homeomorphism is a *simplicial homeomorphism* from some simplicial complex K to another L .

⁶In lemma 5, we will show $\underline{P}_{\text{diff}, \gamma}$ is equal to $P_{\text{diff}, \gamma}$ in [1].

The inner product on this space is defined as follows: For $\underline{\eta}\Psi, \underline{\eta}\Phi \in \text{Im}\underline{\eta}$,

$$\langle \underline{\eta}\Psi, \underline{\eta}\Phi \rangle := (\underline{\eta}\Psi | \Phi). \quad (11)$$

The Cauchy completion of $\underline{\text{Cyl}}_{\text{diff}}^*$ with respect to the above inner product we denote by $\underline{\mathcal{H}}_{\text{diff}}$.

The solution to both the Gauss and diffeomorphism constraints is constructed by first defining $\underline{\text{Cyl}}_{\text{diff},G}^* := \underline{\eta}[\underline{\text{Cyl}} \cap \underline{\mathcal{H}}_G] \subset \underline{\text{Cyl}}_{\text{diff}}^*$, and then Cauchy completing to obtain $\underline{\mathcal{H}}_{\text{diff},G} \subset \underline{\mathcal{H}}_{\text{diff}}$.

3 Equivalence of piecewise linear LQG with analytic LQG at the diffeomorphism invariant level

In this section we prove that the diffeomorphism invariant Hilbert space for piecewise linear LQG is naturally isomorphic to the diffeomorphism invariant Hilbert space of standard LQG — provided that for standard LQG one uses a generalized diffeomorphism group such as that advocated by [13].

We begin by proving the key lemma about piecewise linear LQG allowing the equivalence. Essentially it states that Diff equivalence classes of piecewise linear graphs are simply knot classes. Because the analogue of this is also true for piecewise analytic LQG with the choice of diffeomorphism group advocated in [13], one already has a hint of the equivalence of the two theories at the diffeomorphism invariant level. However, to rigorously prove the equivalence, more must be done, and the subsequent part of this section is devoted to this task.

First we give several definitions. Given a simplicial complex K , a *subcomplex* K' is any subset of K such that K' is again a simplicial complex. (Note it is possible for the dimension of K' to be less than that of K). Second, a complex $\tilde{K}$ is said to be a *subdivision* of a complex K if every simplex of $\tilde{K}$ is contained in a simplex of K , and every simplex of K is a union of simplices in $\tilde{K}$. Third, given a simplicial complex K , we define

$$|K| := \cup_{A \in K} A, \quad (12)$$

called the *polyhedron underlying* K . Lastly, we define a piecewise linear graph γ and a 1-complex X to be *compatible* if the image of γ (which we denote by $|\gamma|$) equals $|X|$. By breaking up each edge of a piecewise linear γ into its straight pieces, and taking the set of these line segments and all their endpoints, one obtains the simplest 1-complex compatible with γ . By subdividing the edges further, one obtains other compatible 1-complexes.

We begin by stating a lemma, which is almost identical to (4.4) of [17]:

Lemma 1 (almost (4.4) of Brown [17]). *Let K and L be 3-complexes and let K_1 and L_1 be 1-dimensional subcomplexes of K and L respectively. Suppose $f : |K| \rightarrow |L|$ is a homeomorphism such that $f(|K_1|) = |L_1|$. Then there exists an isotopy $g_t : |K| \rightarrow |L|$ such that*

- (i) $g_0 = f$
- (ii) *there exist subdivisions $\tilde{K}, \tilde{L}, \tilde{K}_1, \tilde{L}_1$ of K, L, K_1, L_1 respectively such that*
 - (a.) $\tilde{K}_1$ and $\tilde{L}_1$ are subcomplexes of $\tilde{K}$ and $\tilde{L}$, respectively,
 - (b.) g_t maps $\tilde{K}_1$ onto $\tilde{L}_1$ for all t , and
 - (c.) g_1 is piecewise linear on $\tilde{K}_1$.

Proof. The proof is exactly the same as that given for (4.4) in [17]; only the statement of the lemma differs. ■

We use the above in proving the following lemma. A generalized version of the Hauptvermutung of algebraic topology for 3-complexes, proved in 1969 [17], plays a key role in the following proof.

Lemma 2. *If $\gamma, \gamma' \in \underline{\Gamma}$ admit a homeomorphism $\xi : M \rightarrow M$ such that $\gamma' = \xi \cdot \gamma$, then there exists $\varphi \in \underline{\text{Diff}}$ such that $\gamma' = \varphi \cdot \gamma$.*

Proof.

First, by theorem 12 in the appendix, there exist simplicial complexes K and L , each triangulating all of $M = \mathbb{R}^3$, such that K contains a one-dimensional subcomplex K_1 compatible with γ , and L contains a one-dimensional subcomplex L_1 compatible with γ' . Because ξ maps γ to γ' , it maps $|K_1|$ to $|L_1|$. We now invoke lemma 1 above; it provides us with subdivisions $\tilde{K}, \tilde{L}, \tilde{K}_1, \tilde{L}_1$ of K, L, K_1, L_1 such that $\tilde{K}_1$ and $\tilde{L}_1$ are subcomplexes of $\tilde{K}$ and $\tilde{L}$, and an isotopy $\xi_t : M \rightarrow M$ such that (i) $\xi_0 = \xi$, (ii) ξ_t maps $\tilde{K}_1$ to $\tilde{L}_1$ for all t , and (iii) ξ_1 is piecewise linear on $\tilde{K}$.

The 3-complexes $\tilde{K}$ and $\tilde{L}$, the subcomplex $\tilde{K}_1$ of $\tilde{K}$, and the homeomorphism ξ_1 now satisfy the hypotheses of theorem (4.8) of [17], which implies the existence of an isotopy $\varphi_t : M \rightarrow M$, such that (i) $\varphi_0 = \xi_1$, (ii) φ_1 is piecewise linear, and (iii) $\varphi_t|_{|\tilde{K}_1|} = \xi_1|_{|\tilde{K}_1|}$ for all t .

Now, as already noted, ξ_t maps $\tilde{K}_1$ as a 1-complex onto $\tilde{L}_1$ as a 1-complex for all t . That is, ξ_t maps each simplex of $\tilde{K}_1$ to a corresponding simplex of $\tilde{L}_1$ in an onto fashion; this mapping is furthermore 1-1 from the injectivity of ξ_t . Now, because $\tilde{K}_1$ is a subdivision of K_1 , and K_1 is compatible with γ , $\tilde{K}_1$ is also compatible with γ , so that each edge of γ is a union of simplices in $\tilde{K}_1$. Likewise, each edge of γ' is a union of simplices in $\tilde{L}_1$. It follows that, for all t , ξ_t maps each edge of γ onto a corresponding edge of γ' in a 1-1 and onto fashion. The continuity of ξ_t in t ensures that ξ_t always maps each edge of γ to the *same* edge of γ' for all t . Furthermore, recall that $\xi_0 = \xi$ maps the orientation of each edge in γ correctly into the orientation of the corresponding edge in γ' ; the continuity of ξ_t in t ensures that ξ_t does the same for all t . Thus, for all t , ξ_t maps γ onto γ' as a graph. This is in particular true for ξ_1 ; property (iii) of φ_t then implies that this is also true for φ_t for all t . $\varphi := \varphi_1$ thus provides a piecewise linear homeomorphism, i.e., an element of $\underline{\text{Diff}}$, mapping γ to γ' , as desired. ■

Let Γ denote the set of piecewise analytic graphs: that is, graphs with a finite number of oriented compact edges, each of which can be subdivided into a finite number of analytic curves.

Definition (probe equivalent). When two graphs $\gamma, \gamma' \in \Gamma$ differ only by a permutation of edges or reversal of edge orientations, we say that γ and γ' are *probe equivalent*. The probe equivalence class of a graph γ we write $[\gamma]_{pr}$.

Let Γ_{pr} denote the set of probe equivalence classes in Γ , as we have let $\underline{\Gamma}_{pr}$ denote the set of probe equivalence classes in $\underline{\Gamma}$. Let Diff denote the class of diffeomorphisms which one wishes to use to solve the diffeomorphism constraint in the piecewise analytic framework. We make the following assumption about Diff :

Assumption. If $\gamma, \gamma' \in \Gamma$ are such that $\gamma' = \xi \cdot \gamma$ for some homeomorphism $\xi : M \rightarrow M$, then there exists $\varphi \in \text{Diff}$ such that $\gamma' = \varphi \cdot \gamma$.

Note that if Diff is chosen to be the stratified analytic diffeomorphisms [14, 15] as advocated in [13], lemma 4 in [13]⁷ assures that this assumption is satisfied. Finally, let $\mathcal{A}$ denote the space of smooth $SU(2)$ connections on M . In defining analytic LQG and its diffeomorphism-invariant Hilbert-space, we follow [1]. In the following, we only introduce the structures necessary to construct the diffeomorphism invariant Hilbert space.

Definition (Piecewise analytic LQG structures).

⁷using the analytic differentiability class

1. Given a graph $\gamma \in \Gamma$, let Cyl_γ denote the set of functions on $\mathcal{A}$ cylindrical with respect to γ (note that for $\gamma \in \underline{\Gamma}$, this is consistent with the prior definition of Cyl_γ). Let $\text{Cyl} := \cup_\gamma \text{Cyl}_\gamma$.
2. Let $\langle \cdot, \cdot \rangle$ denote the standard inner product on Cyl defined using the Haar measure on $SU(2)$ [1, 2]. Let $\mathcal{H}_\gamma$ and $\mathcal{H}$ denote the Cauchy completions of Cyl_γ and Cyl , respectively, with respect to $\langle \cdot, \cdot \rangle$.
3. Let $\mathcal{H}_G$ denote the solution space to the Gauss constraint, consisting as usual in the Cauchy completion of the span of gauge-invariant spin-networks [1, 2].
4. For each $\gamma \in \Gamma$, let $\mathcal{H}'_\gamma$ denote the orthogonal complement, in $\mathcal{H}_\gamma$, of the span of all functions that are constant on at least one edge of γ , so that, as in [1], $\mathcal{H} = \oplus_{[\gamma] \in \Gamma_{\text{pr}}} \mathcal{H}'_\gamma$. Let $\text{Cyl}'_\gamma := \text{Cyl} \cap \mathcal{H}'_\gamma$. (For $\gamma \in \overline{\Gamma}$, these definitions are again consistent with the ones in the piecewise linear framework.)
5. For each $\gamma \in \Gamma$, let Diff_γ be the set of elements in Diff mapping γ back into its probe equivalence class. Let TDiff_γ be the set of elements in Diff that fix γ — i.e., that preserve each edge of γ , including orientation. So defined, Diff_γ is precisely the subset of Diff preserving Cyl'_γ under pull-back, and TDiff_γ is precisely the subset of Diff that acts as the identity on Cyl'_γ under pull-back. Let $GS_\gamma := \text{Diff}_\gamma / \text{TDiff}_\gamma$ where the division is taken with respect to the left-action.
6. For each $\gamma \in \Gamma$, define $P_{\text{diff}, \gamma}$ as the group averaging map from $\mathcal{H}'_\gamma$ to the subspace invariant under GS_γ :

$$P_{\text{diff}, \gamma} \Psi_\gamma := \frac{1}{|GS_\gamma|} \sum_{\varphi \in GS_\gamma} \varphi^* \Psi_\gamma. \quad (13)$$

For each $\Psi_\gamma \in \text{Cyl}'_\gamma$, define $\eta(\Psi_\gamma) \in \text{Cyl}^*$ by

$$(\eta(\Psi_\gamma) | \Phi) := \sum_{\varphi \in \text{Diff} / \text{TDiff}_\gamma} \langle \varphi^* P_{\text{diff}, \gamma} \Psi_\gamma, \Phi \rangle = \frac{1}{|GS_\gamma|} \sum_{\varphi \in \text{Diff} / \text{TDiff}_\gamma} \langle \varphi^* \Psi_\gamma, \Phi \rangle. \quad (14)$$

Piecing these together for all γ defines a map $\eta : \text{Cyl} \rightarrow \text{Cyl}^*$. This is the rigging map for the theory, as defined in [1] (see also [11], and the related [16]).

7. $\text{Cyl}^*_{\text{diff}} := \text{Im} \eta$. For $\eta\Psi, \eta\Phi \in \text{Im} \eta$,

$$\langle \eta\Psi, \eta\Phi \rangle := (\eta\Psi | \Phi). \quad (15)$$

$\mathcal{H}_{\text{diff}}$ is then defined to be the Cauchy completion of $\text{Cyl}^*_{\text{diff}}$ with respect to this inner product. The completion $\mathcal{H}_{\text{diff}, G}$ of the subspace $\text{Cyl}^*_{\text{diff}, G} := \eta[\text{Cyl} \cap \mathcal{H}_G] \subset \mathcal{H}_{\text{diff}}$ is then the solution to both the Gauss and diffeomorphism constraints.

We next prove a few important lemmas which we use.

Lemma 3. *For each $\gamma \in \underline{\Gamma}$, the map*

$$\begin{aligned} F : \quad \underline{GS}_\gamma &\rightarrow GS_\gamma \\ \varphi \circ [\underline{\text{TDiff}}_\gamma] &\mapsto \varphi \circ [\text{TDiff}_\gamma] \end{aligned} \quad (16)$$

is well-defined, and is an isomorphism, showing $\underline{GS}_\gamma \cong GS_\gamma$.

Proof.

F is well-defined

Suppose $\varphi, \xi \in \underline{\text{Diff}}_\gamma$ are such that $\varphi \circ [\underline{\text{TDiff}}_\gamma] = \xi \circ [\underline{\text{TDiff}}_\gamma]$. Then $\varphi^{-1} \circ \xi \in \underline{\text{TDiff}}_\gamma$, whence $\varphi^{-1} \circ \xi \in \text{TDiff}_\gamma$ also, so that $\varphi \circ [\text{TDiff}_\gamma] = \xi \circ [\text{TDiff}_\gamma]$, proving F well-defined.

F is a homomorphism

This is immediate from the definition of multiplication in the two quotient groups.

F is injective

Suppose $\varphi, \xi \in \underline{\text{Diff}}$ are such that $\varphi \circ [\text{TDiff}_\gamma] = \xi \circ [\text{TDiff}_\gamma]$. Then $\varphi^{-1} \circ \xi \in \text{TDiff}_\gamma$. But $\varphi, \xi \in \underline{\text{Diff}}$, so that $\varphi^{-1} \circ \xi \in \underline{\text{Diff}}$, proving furthermore $\varphi^{-1} \circ \xi \in \underline{\text{TDiff}}_\gamma$. It follows $\varphi \circ [\underline{\text{TDiff}}_\gamma] = \xi \circ [\underline{\text{TDiff}}_\gamma]$, proving injectivity.

F is surjective

Let $\varphi \circ [\text{TDiff}_\gamma] \in GS_\gamma$ be given, so that $\varphi \in \text{Diff}_\gamma$. Let $\gamma' := \varphi \cdot \gamma$. As $\varphi \in \text{Diff}_\gamma$, γ' is probe equivalent to γ and so is also in $\underline{\Gamma}$. Furthermore, φ is in particular a homeomorphism, allowing us to invoke lemma 2, so that there exists a $\xi \in \underline{\text{Diff}}$ such that $\gamma' = \xi \cdot \gamma$. This ξ maps γ to γ' , a graph probe equivalent to γ , whence $\xi \in \underline{\text{Diff}}_\gamma$. Furthermore, $(\varphi^{-1} \circ \xi)\gamma = \gamma$, so that $\varphi^{-1} \circ \xi \in \text{TDiff}_\gamma$, whence $\xi \circ [\text{TDiff}_\gamma] = \varphi \circ [\text{TDiff}_\gamma]$. Thus $F(\xi \circ [\underline{\text{TDiff}}_\gamma]) = \varphi \circ [\text{TDiff}_\gamma]$, proving surjectivity. $\blacksquare$

Lemma 4.

1. Given $\gamma \in \Gamma$, there exists $\varphi \in \text{Diff}$ such that $\varphi \cdot \gamma \in \underline{\Gamma}$.
2. Given $\Psi \in \text{Cyl}$, there exists $\tilde{\varphi} \in \text{Diff}$ such that $\tilde{\varphi}^* \Psi \in \underline{\text{Cyl}}$.

Proof.

Proof of (1.):

Let α be any element of $\underline{\Gamma}$ with the same knot-class as γ (it easy to see that one can construct an element of $\underline{\Gamma}$ with any desired knot-class), and choose the ordering and orientation of the edges of α such that $\alpha = \xi \cdot \gamma$ for some homeomorphism $\xi : M \rightarrow M$. Assumption 3 implies there exists $\varphi \in \text{Diff}$ such that $\alpha = \varphi \cdot \gamma$.

Proof of (2.):

As $\Psi \in \text{Cyl}$, $\Psi \in \text{Cyl}_\gamma$ for some $\gamma \in \Gamma$. From part (1.), there exists $\varphi \in \text{Diff}$ such that $\varphi \cdot \gamma \in \underline{\Gamma}$, so that $(\varphi^{-1})^* \Psi \in \underline{\text{Cyl}}$. $\blacksquare$

Because $\underline{\text{Cyl}} \subset \text{Cyl}$, we have a natural map $\mathcal{J} : \text{Cyl}^* \rightarrow \underline{\text{Cyl}}^*$ defined by

$$(\mathcal{J}\Psi|\Phi) := (\Psi|\Phi) \quad (17)$$

for all $\Psi \in \text{Cyl}^*$ and $\Phi \in \underline{\text{Cyl}}$.

Lemma 5.

1. For $\gamma \in \underline{\Gamma}$, $P_{\text{diff}, \gamma} = \underline{P}_{\text{diff}, \gamma}$.
2. For $\Psi \in \underline{\text{Cyl}}$, $\mathcal{J}\eta\Psi = \underline{\eta}\Psi$.

Proof.

Proof of (1.):

We use the isomorphism F from lemma 3. It is immediate from its definition that, for $\Psi \in \text{Cyl}'_\gamma$ and $\xi \in \underline{\text{Diff}}_\gamma / \underline{\text{TDiff}}_\gamma$, $F(\xi)^* \Psi = \xi^* \Psi$. Using F and this fact,

$$P_{\text{diff}, \gamma} \Psi := \frac{1}{|\underline{GS}_\gamma|} \sum_{\varphi \in \underline{GS}_\gamma} \varphi^* \Psi = \frac{1}{|\underline{GS}_\gamma|} \sum_{\varphi \in \underline{GS}_\gamma} (F\varphi)^* \Psi = \frac{1}{|\underline{GS}_\gamma|} \sum_{\varphi \in \underline{GS}_\gamma} \varphi^* \Psi = \underline{P}_{\text{diff}, \gamma} \Psi. \quad (18)$$

Proof of (2.):

Using the linearity of $\mathcal{J}$, η and η' , without loss of generality, assume $\Psi \in \text{Cyl}'_\gamma$ for some $\gamma \in \underline{\Gamma}$. Suppose $\gamma' \in \underline{\Gamma}$ and $\Theta \in \text{Cyl}'_{\gamma'}$ are given.

Case 1: There exists no $\varphi_o \in \text{Diff}$ such that $\varphi_o \cdot \gamma' = \gamma$.

Then from (14), $(\mathcal{I}\eta\Psi \mid \Theta) = 0$. But from (9), $(\underline{\eta}\Psi \mid \Theta) = 0$ as well, so that $(\mathcal{I}\eta\Psi \mid \Theta) = (\underline{\eta}\Psi \mid \Theta) = 0$.

Case 2: There exists $\varphi_o \in \text{Diff}$ such that $\varphi_o \cdot \gamma' = \gamma$.

Then, from lemma 2, there exists $\underline{\varphi}_o \in \underline{\text{Diff}}$ such that $\underline{\varphi}_o \cdot \gamma' = \gamma$. Using the orthogonality of the spaces $\mathcal{H}'_\gamma$, the middle expression in (14) reduces to

$$(\mathcal{I}\eta\Psi \mid \Theta) = (\eta\Psi \mid \Theta) = \langle \underline{\varphi}_o^* P_{\text{diff}, \gamma} \Psi, \Theta \rangle \quad (19)$$

Using part (1.) of this lemma, and the same orthogonality of the spaces $\mathcal{H}'_\gamma$ to simplify the expression for $(\underline{\eta}\Psi \mid \Theta)$, we also have

$$(\mathcal{I}\eta\Psi \mid \Theta) = \langle \underline{\varphi}_o^* P_{\text{diff}, \gamma} \Psi, \Theta \rangle = (\underline{\eta}\Psi \mid \Theta) \quad (20)$$

Thus $(\mathcal{I}\eta\Psi \mid \Theta) = (\underline{\eta}\Psi \mid \Theta)$ for all $\Theta \in \text{Cyl}'_{\gamma'}$, $\gamma' \in \underline{\Gamma}$, so that $\mathcal{I}\eta\Psi = \underline{\eta}\Psi$. ■

Theorem 6. $\mathcal{I}$ maps $\text{Cyl}_{\text{diff}}^*$ onto $\underline{\text{Cyl}}_{\text{diff}}^*$. Furthermore, $\mathcal{I}|_{\text{Cyl}_{\text{diff}}^*} : \text{Cyl}_{\text{diff}}^* \rightarrow \underline{\text{Cyl}}_{\text{diff}}^*$ is a unitary isomorphism.

Proof.

Proof that $\mathcal{I}[\text{Cyl}_{\text{diff}}^*] = \underline{\text{Cyl}}_{\text{diff}}^*$:

$\subseteq$:

Let $\eta\Psi \in \text{Cyl}_{\text{diff}}^*$ be given, so that $\Psi \in \text{Cyl}$. By lemma 4, $\exists \xi \in \text{Diff}$ s.t. $\xi^*\Psi \in \text{Cyl}$. Using the Diff invariance of η and part (2.) of lemma 5, $\mathcal{I}\eta\Psi = \mathcal{I}\eta(\xi^*\Psi) = \underline{\eta}(\xi^*\Psi)$, which is in $\underline{\text{Cyl}}_{\text{diff}}^*$.

$\supseteq$:

Let $\underline{\eta}\Psi \in \underline{\text{Cyl}}_{\text{diff}}^*$ be given, so $\Psi \in \underline{\text{Cyl}}$. Then $\eta\Psi \in \text{Cyl}_{\text{diff}}^*$, and by lemma 5, $\mathcal{I}\eta\Psi = \underline{\eta}\Psi$, so that $\underline{\eta}\Psi \in \mathcal{I}[\text{Cyl}_{\text{diff}}^*]$.

Proof that $\mathcal{I}|_{\text{Cyl}_{\text{diff}}^*}$ is injective:

Suppose $\eta\Psi, \eta\Phi \in \text{Cyl}_{\text{diff}}^*$ are such that $\mathcal{I}\eta\Psi = \mathcal{I}\eta\Phi$. Let $\Theta \in \text{Cyl}$ be given. By lemma 4, there exists $\xi \in \text{Diff}$ such that $\xi^*\Theta \in \underline{\text{Cyl}}$. Using the Diff invariance of η ,

$$(\eta\Psi \mid \Theta) = (\eta\Psi \mid \xi^*\Theta) = (\mathcal{I}\eta\Psi \mid \xi^*\Theta) = (\mathcal{I}\eta\Phi \mid \xi^*\Theta) = (\eta\Phi \mid \xi^*\Theta) = (\eta\Phi \mid \Theta)$$

for all $\Theta \in \text{Cyl}$, whence $\eta\Psi = \eta\Phi$.

Proof that $\mathcal{I}|_{\text{Cyl}_{\text{diff}}^*}$ is isometric and hence unitary:

Let $\eta\Psi, \eta\Phi \in \text{Cyl}_{\text{diff}}^*$ be given, so that $\Psi, \Phi \in \text{Cyl}$. Using lemma 4, there exists φ and ξ in Diff such that $\varphi^*\Psi, \xi^*\Phi \in \underline{\text{Cyl}}$. Using the Diff invariance of η and part (2.) of lemma 5, we have

$$\begin{aligned} \langle \mathcal{I}\eta\Psi, \mathcal{I}\eta\Phi \rangle &= \langle \mathcal{I}\eta(\varphi^*\Psi), \mathcal{I}\eta(\xi^*\Phi) \rangle = \langle \underline{\eta}(\varphi^*\Psi), \underline{\eta}(\xi^*\Phi) \rangle \\ &:= \langle \underline{\eta}(\varphi^*\Psi) \mid \xi^*\Phi \rangle = \langle \mathcal{I}\eta(\varphi^*\Psi) \mid \xi^*\Phi \rangle = \langle \eta(\varphi^*\Psi) \mid \xi^*\Phi \rangle \\ &= \langle \eta\Psi \mid \Phi \rangle = \langle \eta\Psi, \eta\Phi \rangle, \end{aligned}$$

The above theorem implies ■

Corollary 7. $\mathcal{H}_{\text{diff}}$ and $\underline{\mathcal{H}}_{\text{diff}}$ are isomorphic as Hilbert spaces.

It is then easy to extend the equivalence to the solution spaces solving both the diffeomorphism and Gauss constraints:

Corollary 8. $\mathcal{J}|_{\text{Cyl}_{\text{diff},G}^*} : \underline{\text{Cyl}}_{\text{diff},G}^* \rightarrow \text{Cyl}_{\text{diff},G}^*$ is a unitary isomorphism, so that $\mathcal{H}_{\text{diff},G}$ and $\underline{\mathcal{H}}_{\text{diff},G}$ are isomorphic as Hilbert spaces.

Proof. From the injectivity of $\mathcal{J}|_{\text{Cyl}_{\text{diff}}^*}$, we know $\mathcal{J}|_{\text{Cyl}_{\text{diff},G}^*}$ is injective. It thus remains only to prove that $\mathcal{J}$ maps $\text{Cyl}_{\text{diff},G}^*$ onto $\underline{\text{Cyl}}_{\text{diff},G}^*$, i.e., $\mathcal{J}[\text{Cyl}_{\text{diff},G}^*] = \underline{\text{Cyl}}_{\text{diff},G}^*$.

($\subseteq$):

Let $\eta\Psi \in \text{Cyl}_{\text{diff},G}^*$ be given, so that $\Psi \in \text{Cyl} \cap \mathcal{H}_G$, and in particular $\Psi \in \text{Cyl}_\gamma$ for some $\gamma \in \Gamma$. By part (1.) of lemma 4, there exists $\varphi \in \text{Diff}$ such that $\varphi \cdot \gamma \in \underline{\Gamma}$. Then $(\varphi^{-1})^*\Psi \in \underline{\text{Cyl}} \cap \underline{\mathcal{H}}_G$, and we have $\mathcal{J}\eta\Psi = \mathcal{J}\eta(\varphi^{-1})^*\Psi = \underline{\eta}(\varphi^{-1})^*\Psi$, where lemma 2 was used in the second step. Thus $\mathcal{J}\eta\Psi \in \underline{\text{Cyl}}_{\text{diff},G}^*$.

($\supseteq$):

Let $\underline{\eta}\Psi \in \underline{\text{Cyl}}_{\text{diff},G}^*$ be given, so that $\Psi \in \underline{\text{Cyl}} \cap \underline{\mathcal{H}}_G \subset \text{Cyl} \cap \mathcal{H}_G$. By lemma 2, $\mathcal{J}\eta\Psi = \underline{\eta}\Psi$, so that $\underline{\eta}\Psi \in \mathcal{J}[\text{Cyl}_{\text{diff},G}^*]$. $\blacksquare$

4 Equivalence of diffeomorphism invariant operators, and equivalence of dynamics

When constructing operators in pLQG, we propose one quantize in exactly the same way as in standard LQG, except that only piecewise linear edges should be used. For operators preserving Cyl , this general statement can be made precise as follows. Given an operator $\hat{O}_\omega$ in standard LQG, a corresponding operator is defined in pLQG iff $\hat{O}_\omega$ preserves $\underline{\text{Cyl}}$, and in this case one defines the corresponding operator $\hat{O}_{pl}$ in pLQG to be $\hat{O}_\omega|_{\underline{\text{Cyl}}}$. An immediate consequence of this definition is

$$\hat{O}_{pl}^* \circ \mathcal{J} = \mathcal{J} \circ \hat{O}_\omega^*. \quad (21)$$

Next, we call an operator “diffeomorphism invariant” if it is invariant under the group of generalized diffeomorphisms in the relevant framework. If $\hat{O}_\omega$ preserves Cyl and is diffeomorphism invariant, then it must be graph preserving⁸ and hence also preserve $\underline{\text{Cyl}}$, so that there is a corresponding piecewise linear operator $\hat{O}_{pl}$. Because $\underline{\text{Diff}} \subset \text{Diff}$, the Diff-invariance of $\hat{O}_\omega$ also implies the $\underline{\text{Diff}}$ -invariance of $\hat{O}_{pl}$, so that $\hat{O}_{pl}$ is diffeomorphism invariant. These observations, along with (21) allow us to state the following

Proposition 9. *Given any diffeomorphism-invariant operator $\hat{O}_\omega$ preserving Cyl in standard LQG, then $\hat{O}_\omega$ also preserves $\underline{\text{Cyl}}$. The corresponding piecewise linear operator $\hat{O}_{pl}$ is also diffeomorphism invariant, and $\hat{O}_{pl}$ and $\hat{O}_\omega$ are mapped into each other by the isomorphism $\mathcal{J}|_{\text{Cyl}_{\text{diff}}^*}$, that is,*

$$\hat{O}_{pl}^* \circ \mathcal{J} = \mathcal{J} \circ \hat{O}_\omega^*. \quad (22)$$

⁸This can be seen as follows. Suppose $\hat{O}$ is Diff-invariant and preserves Cyl . Let $\Psi \in \text{Cyl}_\alpha$ be given for some α . As $\hat{O}$ preserves Cyl , $\hat{O}\Psi \in \text{Cyl}_\beta$ for some β . From Diff-invariance, we have that for all $\varphi \in \text{Diff}_\alpha$ (recall Diff_α is the subgroup of Diff preserving α), $\hat{O}\Psi = U_\varphi \hat{O} U_{\varphi^{-1}} \Psi = U_\varphi \hat{O} \Psi$, so that $\hat{O}\Psi \in \text{Cyl}_{\varphi \cdot \beta}$ for all $\varphi \in \text{Diff}_\alpha$. Thus $\hat{O}\Psi \in \cup_{\varphi \in \text{Diff}_\alpha} \text{Cyl}_{\varphi \cdot \beta}$. But given any $\gamma, \gamma' \in \Gamma$, $\text{Cyl}_\gamma \cap \text{Cyl}_{\gamma'} = \text{Cyl}_{\gamma \cap \gamma'}$, so that $\hat{O}\Psi \in \text{Cyl}_{\cap_{\varphi \in \text{Diff}_\alpha} \varphi \cdot \beta}$. The only edges of β that survive in $\cap_{\varphi \in \text{Diff}_\alpha} \varphi \cdot \beta$ are those that are also edges of α , whence in fact $\hat{O}\Psi \in \text{Cyl}_\alpha$, showing $\hat{O}$ is graph preserving.

Note that, though this proposition seems quite general, in fact assuming $\hat{O}_\omega$ is both well defined on Cyl and diffeomorphism invariant is a relatively restrictive assumption: it already constrains the applicability of the result to graph preserving operators. The master constraint operator [18], for example, though diffeomorphism invariant, is not graph preserving. This is possible because the master constraint is not well-defined on Cyl , but rather must be directly defined on $\text{Cyl}_{\text{diff}}^*$. We will later discuss the master constraint, after we have discussed the Hamiltonian constraint.

The Hamiltonian constraint [19] is rather unique because it has as its domain $\text{Cyl}_{\text{diff}}^*$, but does not map $\text{Cyl}_{\text{diff}}^*$ back into itself. It is defined as follows. For each lapse N , each $\epsilon > 0$ and each graph γ , one defines a regulated operator $\hat{H}(N)_{\gamma,\epsilon}$ on $\mathcal{H}'_\gamma$. Piecing these together for all γ gives, for each ϵ , an operator $\hat{H}(N)_\epsilon$ on the kinematical Hilbert space $\mathcal{H}$. The dual $\hat{H}(N)_\epsilon^*$ then acts on Cyl^* . For any ξ in $\text{Cyl}_{\text{diff}}^* \subset \text{Cyl}^*$, the limit $\lim_{\epsilon \rightarrow 0} \hat{H}(N)_\epsilon^* \xi$ becomes trivial, allowing us to define

$$\hat{H}(N)\xi := \lim_{\epsilon \rightarrow 0} \hat{H}(N)_\epsilon^* \xi. \quad (23)$$

$\hat{H}(N)$ is thus well-defined on $\text{Cyl}_{\text{diff}}^*$. It is also diffeomorphism covariant: $(U_\varphi^{-1})^* \circ \hat{H}(N) \circ U_\varphi^* = \hat{H}(\varphi^* N)$ for all $\varphi \in \text{Diff}$, where U_φ denotes the unitary action of φ on $\mathcal{H}$ via pullback. However, for general lapse N , $\hat{H}(N)$ maps $\text{Cyl}_{\text{diff}}^*$ out of itself due to $\hat{H}(N)$ not being diffeomorphism *invariant*. One can nevertheless define the solution to the Hamiltonian constraint to be simply the common kernel of the operators $\hat{H}(N)$ for all lapse N .

This construction can be repeated in the obvious way for pLQG: one need only ensure that the loops added by the regulated $\hat{H}(N)_{\gamma,\epsilon}$ are chosen to be piecewise linear. We do this, and then for $\gamma \in \underline{\Gamma}$, define $\hat{H}(N)_{\gamma,\epsilon} := \hat{H}(N)_{\gamma,\epsilon}|_{\underline{\text{Cyl}}}$. A construction exactly parallel to that above then goes through, giving us a family of operators $\hat{H}(N)$, defined on $\underline{\text{Cyl}}_{\text{diff}}^*$, and diffeomorphism covariant with respect to $\underline{\text{Diff}}$, which nevertheless generically map $\underline{\text{Cyl}}_{\text{diff}}^*$ out of itself.

Let $\ker \hat{H}$ denote the common kernel of the operators $\hat{H}(N)$ for all N , and let $\ker \hat{H}$ denote the common kernel of the operators $\hat{H}(N)$ for all N . We have the following result:

Proposition 10. $\mathcal{J}|_{\ker \hat{H}}$ provides a unitary isomorphism from $\ker \hat{H}$ onto $\ker \hat{H}$.

Proof.

We first note that for $\Psi \in \text{Cyl}_{\text{diff}}^*$, $\Phi \in \underline{\text{Cyl}}$, and any lapse N , the following relation holds:

$$\begin{aligned} (\hat{H}(N)\mathcal{J}\Psi|\Phi) &:= \lim_{\epsilon \rightarrow 0} (\mathcal{J}\Psi|\hat{H}(N)_\epsilon \Phi) \\ &= \lim_{\epsilon \rightarrow 0} (\Psi|\hat{H}(N)_\epsilon \Phi) = \lim_{\epsilon \rightarrow 0} (\Psi|\hat{H}(N)_\epsilon \Phi) \\ &= (\hat{H}(N)\Psi|\Phi). \end{aligned} \quad (24)$$

From this we immediately see that if $\Psi \in \ker \hat{H}$, so that $\hat{H}(N)\Psi = 0$ for all N , then $\hat{H}(N)\mathcal{J}\Psi = 0$ for all N , so that $\mathcal{J}\Psi \in \ker \hat{H}$, whence $\mathcal{J}[\ker \hat{H}] \subset \ker \hat{H}$.

To prove the converse, let $\Theta \in \ker \hat{H}$ be given. As $\underline{\text{Cyl}}_{\text{diff}}^*$ is defined to be the domain of the $\hat{H}(N)$, $\Theta \in \underline{\text{Cyl}}_{\text{diff}}^*$; using the onto-ness of $\mathcal{J}|_{\underline{\text{Cyl}}_{\text{diff}}^*} : \underline{\text{Cyl}}_{\text{diff}}^* \rightarrow \underline{\text{Cyl}}_{\text{diff}}^*$, there exists $\Psi \in \text{Cyl}_{\text{diff}}^*$ such that $\Theta = \mathcal{J}\Psi$. Next, let N be given, and let $\Phi \in \underline{\text{Cyl}}$ be given. By lemma 4, there exists $\varphi \in \text{Diff}$ such that $\varphi^* \Phi \in \underline{\text{Cyl}}$. Using the Diff covariance of $\hat{H}(N)$ and then the Diff invariance of $(\Psi|$,

$$\begin{aligned} (\hat{H}(N)\Psi|\Phi) &= ((U_\varphi^{-1})^* \circ \hat{H}((\varphi^{-1})^* N) \circ (U_\varphi)^* \Psi|\Phi) = (\hat{H}((\varphi^{-1})^* N)\Psi|U_\varphi^{-1}|\Phi) \\ &= (\hat{H}((\varphi^{-1})^* N)\Psi|\varphi^* \Phi). \end{aligned} \quad (25)$$

Applying relation (24) to $\varphi^* \Phi$ and $(\varphi^{-1})^* N$, and then using the fact that $\Theta = \mathcal{J}\Psi$ is in $\ker \hat{H}$, the last line above is seen to be zero. Thus $(\hat{H}(N)\Psi|\Phi) = 0$ for all $\Phi \in \underline{\text{Cyl}}$ and all lapse N , proving

$\Psi \in \ker \hat{H}$, so that $\Theta \in \mathcal{J}[\ker \hat{H}]$. This proves the containment $\ker \hat{H} \subset \mathcal{J}[\ker \hat{H}]$, completing the proof that $\ker \hat{H} = \mathcal{J}[\ker \hat{H}]$.

As already shown in theorem 6, $\mathcal{J}$ is injective and unitary on $\text{Cyl}_{\text{diff}}^*$, so that it is also injective and unitary on $\ker \hat{H}$. Thus $\mathcal{J}|_{\ker \hat{H}} : \ker \hat{H} \rightarrow \ker \hat{H}$ provides a unitary isomorphism between $\ker \hat{H}$ and $\ker \hat{H}$. ■

Finally, the physical Hilbert space of solutions to the diffeomorphism, Gauss, and Hamiltonian constraint in LQG and pLQG are $\mathcal{H}_{\text{Phys}} := \overline{\text{Cyl}_{\text{diff},G}^* \cap \ker \hat{H}}$ and $\underline{\mathcal{H}}_{\text{Phys}} := \overline{\text{Cyl}_{\text{diff},G}^* \cap \ker \hat{H}}$, respectively, where the closure denotes Cauchy completion. As the isomorphism $\mathcal{J}|_{\text{Cyl}_{\text{diff}}^*}$ maps the inner product on $\text{Cyl}_{\text{diff}}^*$ onto that on $\underline{\text{Cyl}_{\text{diff}}^*}$, maps $\text{Cyl}_{\text{diff},G}^*$ onto $\underline{\text{Cyl}_{\text{diff},G}^*}$, and maps $\ker \hat{H}$ onto $\ker \hat{H}$, it is immediate that $\mathcal{J}$ provides a unitary isomorphism between these physical Hilbert spaces.

We now come to the master constraint. Let us review its construction in standard LQG from [18]. First, given a spatial point $v \in M$, let $N_v(x) := \delta_{v,x}$, a particular singular choice of lapse. The corresponding Hamiltonian constraint operator $\hat{H}_v := \hat{H}(N_v)$ is nevertheless well defined [19], as is perhaps not surprising given the discreteness of LQG. We next recall the *generalized spin-network* functions T_σ , where σ denotes the triple $(\gamma, \vec{j}, \vec{T})$ of a graph $\gamma \in \Gamma$, an assignment of a spin to each edge, and an assignment of a tensor among representations to each vertex [1, 2]. We require that all spin labels be non-trivial. Furthermore, as in, e.g., [1], for each possible set of representations incident at a vertex, we fixed a basis of the tensor space among the representations. Let $\mathcal{S}$ denote the space of all such triples $(\gamma, \vec{j}, \vec{T})$. $\{T_\sigma\}_{\sigma \in \mathcal{S}}$ forms an orthonormal basis of Cyl and hence $\mathcal{H}$. Furthermore, Diff acts on $\mathcal{S}$, so that we may consider the Diff -equivalence class of an element $\sigma \in \mathcal{S}$, which we denote $[\sigma]_{\text{Diff}}$. With these definitions made, we define a quadratic form $Q_M : \text{Cyl}_{\text{diff}}^* \times \text{Cyl}_{\text{diff}}^* \rightarrow \mathbb{C}$ by

$$Q_M(\Phi, \Psi) := \sum_{[\sigma]_{\text{Diff}}} \eta_{[\sigma]_{\text{Diff}}} \sum_{v \in V(\gamma(\sigma))} \overline{(\hat{H}_v \Phi | T_\sigma)} (\hat{H}_v \Psi | T_\sigma). \quad (26)$$

where $\eta_{[\sigma]_{\text{Diff}}} := 1/|GS_{\gamma(\sigma)}|$ are the coefficients appearing in the last expression in (14) for the diffeomorphism constraint rigging map, and where $V(\gamma(\sigma))$ denotes the set of vertices in $\gamma(\sigma)$.⁹ $Q_M(\cdot, \cdot)$ then determines the master constraint $\hat{M}$ uniquely via [18]

$$\hat{M}\Phi := \sum_{x \in I} Q_M(B_x, \Phi) B_x \quad (27)$$

where $\{B_x\}_{x \in I}$ is any orthonormal basis of $\text{Cyl}_{\text{diff}}^*$.

A construction parallel to the above goes through in the pLQG case. Let $\underline{\mathcal{S}}$ denote the set of generalized spin-network labels $\sigma = (\gamma, \vec{j}, \vec{T})$ such that $\gamma \in \underline{\Gamma}$. Then $\underline{\text{Diff}}$ acts on $\underline{\mathcal{S}}$, so that for each $\sigma \in \underline{\text{Diff}}$, one can define an equivalence class $[\sigma]_{\underline{\text{Diff}}}$. The quadratic form for the piecewise linear framework is then

$$Q_{\underline{M}}(\Phi, \Psi) := \sum_{[\sigma]_{\underline{\text{Diff}}}} \eta_{[\sigma]_{\underline{\text{Diff}}}} \sum_{v \in V(\gamma(\sigma))} \overline{(\hat{H}_v \Phi | T_\sigma)} (\hat{H}_v \Psi | T_\sigma) \quad (28)$$

where $\Phi, \Psi \in \underline{\text{Cyl}_{\text{diff}}^*}$, and where $\eta_{[\sigma]_{\underline{\text{Diff}}}} = 1/|\underline{GS}_{\gamma(\sigma)}|$ are the coefficients in the pLQG rigging map (9). The master constraint is then

$$\hat{M}\Phi := \sum_{x \in I} Q_{\underline{M}}(\underline{B}_x, \Phi) \underline{B}_x \quad (29)$$

⁹In [18], $\eta_{[\sigma]_{\text{Diff}}}$ are a set of constants parametrizing an ambiguity in the definition of the rigging map discussed in the original work [11]. Here, as earlier in this paper, we are taking a natural resolution to this ambiguity suggested in [1], leading to the specific values of $\eta_{[\sigma]_{\text{Diff}}}$ given above.

where $\{\underline{B}_x\}_{x \in I}$ is any orthonormal basis of $\underline{\text{Cyl}}_{\text{diff}}^*$.

Proposition 11. $\hat{M}$ is mapped into $\hat{\underline{M}}$ by the isomorphism $\mathcal{J}|_{\text{Cyl}_{\text{diff}}^*}$.

Proof. In each case the master constraint is determined from the quadratic form and inner product on diffeomorphism invariant states in the same way. To prove equivalence of the master constraints, it is thus sufficient to prove equivalence of the quadratic forms; that is, we want to show $Q_{\underline{M}}(\mathcal{J}\Phi, \mathcal{J}\Psi) = Q_M(\Phi, \Psi)$ for all $\Phi, \Psi \in \text{Cyl}_{\text{diff}}^*$:

$$\begin{aligned}
Q_{\underline{M}}(\mathcal{J}\Phi, \mathcal{J}\Psi) &:= \sum_{[\sigma]_{\underline{\text{Diff}}}} \eta_{[\sigma]_{\underline{\text{Diff}}}} \sum_{v \in V(\gamma(\sigma))} \overline{(\hat{\underline{H}}_v \mathcal{J}\Phi | T_\sigma)} (\hat{\underline{H}}_v \mathcal{J}\Psi | T_\sigma) \\
&= \sum_{[\sigma]_{\underline{\text{Diff}}}} \eta_{[\sigma]_{\underline{\text{Diff}}}} \sum_{v \in V(\gamma(\sigma))} \lim_{\epsilon, \epsilon' \rightarrow \infty} \overline{(\mathcal{J}\Phi | \hat{\underline{H}}_{v, \epsilon} T_\sigma)} (\mathcal{J}\Psi | \hat{\underline{H}}_{v, \epsilon} T_\sigma) \\
&= \sum_{[\sigma]_{\underline{\text{Diff}}}} \eta_{[\sigma]_{\underline{\text{Diff}}}} \sum_{v \in V(\gamma(\sigma))} \lim_{\epsilon, \epsilon' \rightarrow \infty} \overline{(\Phi | \hat{H}_{v, \epsilon} T_\sigma)} (\Psi | \hat{H}_{v, \epsilon} T_\sigma) \\
&= \sum_{[\sigma]_{\underline{\text{Diff}}}} \eta_{[\sigma]_{\underline{\text{Diff}}}} \sum_{v \in V(\gamma(\sigma))} \overline{(\hat{H}_v \Phi | T_\sigma)} (\hat{H}_v \Psi | T_\sigma)
\end{aligned} \tag{30}$$

where, in the third equality, we have used the definition of $\mathcal{J}$ and that $\hat{\underline{H}}_{v, \epsilon} = \hat{H}_{v, \epsilon}|_{\underline{\text{Cyl}}}$. Now, the outer sum in (30) is over $[\sigma]_{\underline{\text{Diff}}} \in \underline{\mathcal{S}}/\underline{\text{Diff}}$. Define $J : \underline{\mathcal{S}}/\underline{\text{Diff}} \rightarrow \mathcal{S}/\text{Diff}$ by $[\sigma]_{\underline{\text{Diff}}} \mapsto [\sigma]_{\text{Diff}}$. J is well-defined due to $\underline{\text{Diff}} \subset \text{Diff}$. Using lemma 2, one shows that it is 1-1, and using lemma 4 one sees that it is onto. (Details: exercise for the reader.) Furthermore, as $\sigma \in \underline{\mathcal{S}}$, $\gamma(\sigma) \in \underline{\mathcal{I}}$, so that from lemma 3, $|\underline{GS}_{\gamma(\sigma)}| = |GS_{\gamma(\sigma)}|$, and we have $\eta_{[\sigma]_{\underline{\text{Diff}}}} = \eta_{[\sigma]_{\text{Diff}}}$. Using the isomorphism J to replace $[\sigma]_{\underline{\text{Diff}}}$ with $[\sigma]_{\text{Diff}}$ in (30), we obtain

$$Q_{\underline{M}}(\mathcal{J}\Phi, \mathcal{J}\Psi) = Q_M(\Phi, \Psi). \tag{31}$$

Lastly, because $\mathcal{J}|_{\text{Cyl}_{\text{diff}}^*}$ maps the master constraint $\hat{M}$ onto $\hat{\underline{M}}$, and $\text{Cyl}_{\text{diff}, G}^*$ onto $\underline{\text{Cyl}}_{\text{diff}, G}^*$, $\mathcal{J}|_{\text{Cyl}_{\text{diff}, G}^*}$ will map $\hat{M}|_{\text{Cyl}_{\text{diff}, G}^*}$ onto $\hat{\underline{M}}|_{\underline{\text{Cyl}}_{\text{diff}, G}^*}$, so that the master constraint dynamics are also equivalent after solving both the diffeomorphism and Gauss constraints.¹⁰

The above results show that not only are the diffeomorphism invariant Hilbert spaces in LQG and pLQG unitarily isomorphic, but the dynamics (whether defined with Master or Hamiltonian constraint) are isomorphic as well, so that the two frameworks are truly equivalent.

5 Exact embedding of LQC into piecewise linear LQG

In the paper [5], an embedding of LQC into the space $\underline{\text{Cyl}}^*$ was constructed. In that context, the space $\underline{\text{Cyl}}^*$ was unnatural as a distributional space in the sense that it was the dual of a test function space that is not dense in the traditional kinematical Hilbert space $\mathcal{H}$ of LQG. A possible physical meaning for $\underline{\text{Cyl}}^*$ was suggested in [5], but this did not solve the fact that it was not clear how to use $\underline{\text{Cyl}}^*$ for the next step in the program of [4, 5]. Specifically, the next step was to group average the kinematical embeddings to obtain embeddings into LQG at the diffeomorphism invariant level. Note one *must* construct embeddings into LQG at the diffeomorphism invariant level if one hopes to

¹⁰In [18], the master constraint is in fact constructed directly on $\text{Cyl}_{\text{diff}, G}^*$.

exactly relate the Hamiltonian constraints in LQC and LQG in any way, as the latter is defined only on diffeomorphism invariant states. To accomplish the construction of the diffeomorphism invariant embeddings, two issues needed to be addressed [5]:

1. The group of piecewise analytic diffeomorphisms did not even act on $\underline{\text{Cyl}}^*$, so that one could not even write down a formal expression for group averaging the kinematical embeddings.
2. Once one is able to write down a formal group averaging, one would need to regulate the integral over diffeomorphisms in some way.

It is in this first step that the use of $\underline{\text{Cyl}}^*$ seemed to prevent further progress.

In the construction of pLQG, $\underline{\text{Cyl}}^*$ also appears, but this time *as the space of distributional states for a completely parallel framework for loop quantum gravity, which, as was proven above, is equivalent to the standard one at the diffeomorphism-invariant level*. Furthermore, the space of ‘piecewise linear generalized diffeomorphisms’ acts on $\underline{\text{Cyl}}^*$, so that one can now formally write down the group averaging of the embeddings, providing an expression for the embedding into the space of diffeomorphism invariant states. Because of the isomorphism between pLQG and LQG at the diffeomorphism invariant level, this is also a formal expression for the embedding into the space of diffeomorphism invariant states in standard LQG. That is, the first obstruction listed above is gone. Because the embeddings of [5] were a motivation for the present work, we briefly review them here; we then end the section with the new expressions for the diffeomorphism invariant embeddings.

First we recall some necessary structures from loop quantum cosmology (LQG). As in [20], we take the classical configuration space for homogeneous, isotropic cosmology to be the space of homogeneous, isotropic connections, but in the gauge-fixed sense defined in [5]; we denote this space by $\mathcal{A}_S$. By picking a reference connection $\dot{A}_a^i \in \mathcal{A}_S$, all other connections in $\mathcal{A}_S$ are related to $\dot{A}_a^i$ by scaling. Thus if we define $r : \mathbb{R} \rightarrow \mathcal{A}_S \subset \mathcal{A}$ by

$$r : c \mapsto c \dot{A}_a^i, \quad (32)$$

r provides an isomorphism of $\mathbb{R}$ with $\mathcal{A}_S$. States in LQC are then functions on $\mathbb{R} \cong \mathcal{A}_S$. The basic space of ‘nice’ states in LQC (and one of the sources of the unique character of LQC) is the space of *almost periodic functions*; following [20], we denote this Cyl_S . Cyl_S^* is the space of distributional states.

The kinematical and gauge-invariant embeddings of [5] are then defined as follows. The ‘c’ embedding $\iota_c : \text{Cyl}_S^* \rightarrow \underline{\text{Cyl}}^*$ is defined by

$$(\iota_c \psi | \Phi) := (\psi | r^* \Phi). \quad (33)$$

From ι_c , one constructs the ‘b’ embeddings. To remind the reader from [4, 5], the ‘b’ embedding is built using coherent states, the idea being to use the freedom in the choice of coherent states to adapt the embedding to be approximately preserved by the dynamics. In [4, 5], complexifier coherent states are used; in complexifier coherent states, the freedom in choosing the family of coherent states is parametrized by a choice of complexifier [21]. To introduce the complexifiers, first let X_S and X denote the classical phase space of the reduced and full theories, respectively. Then let $C_S : X_S \rightarrow \mathbb{R}^+$, $C : X \rightarrow \mathbb{R}^+$ be any two (pure momentum) complexifiers [21]. Let $\hat{C}_S$ and $\hat{C}$ denote their respective quantizations in the reduced and full quantum theories. For brevity, we give only the final expression for the corresponding ‘b’ embedding $\iota_b : \text{Cyl}_S^* \rightarrow \underline{\text{Cyl}}^*$. It is given by [5]

$$(\iota_b \psi | \Phi) := (\psi | e^{\hat{C}_S} \circ r^* \circ e^{-\hat{C}} | \Phi). \quad (34)$$

The Gauss-gauge invariant versions of these embeddings are $\iota_c^G := P_G^* \circ \iota_c$ and $\iota_b^G := P_G^* \circ \iota_b$, where $P_G : \underline{\text{Cyl}} \rightarrow \underline{\text{Cyl}}$ denotes the projector onto gauge-invariant states. For the motivation behind these definitions and their nice properties, we refer the reader to the original papers [4, 5].

Now we come to the formal expression for the embedding into diffeomorphism invariant states, made possible by the new piecewise LQG framework introduced in this paper. The diffeomorphism invariant embedding $\iota_c^{Diff} : \text{Cyl}_S^* \rightarrow \underline{\text{Cyl}}_{\text{diff}}^*$ has the formal expression

$$\begin{aligned} (\iota_c^{Diff} \psi | \Phi) &:= \left(\int_{\varphi \in \underline{\text{Diff}}} \mathcal{D}\varphi (U_\varphi^* \iota_c^G \psi |) \right) | \Phi \rangle \\ &= \int_{\varphi \in \underline{\text{Diff}}} \mathcal{D}\varphi (\iota_c^G \psi | U_\varphi | \Phi) \rangle \\ &= \int_{\varphi \in \underline{\text{Diff}}} \mathcal{D}\varphi (\psi | r^* \circ P_G \circ U_\varphi | \Phi) \rangle. \end{aligned} \quad (35)$$

The formal expression for the diffeomorphism invariant ‘b’ embedding $\iota_b^{Diff} : \text{Cyl}_S^* \rightarrow \underline{\text{Cyl}}_{\text{diff}}^*$ is then

$$(\iota_b^{Diff} \psi | \Phi) = \int_{\varphi \in \underline{\text{Diff}}} \mathcal{D}\varphi (\psi | e^{\hat{C}_S} \circ r^* \circ e^{-\hat{C}} \circ P_G \circ U_\varphi | \Phi) \rangle. \quad (36)$$

If $\hat{C}$ and $\hat{C}_S$ are gauge and diffeomorphism invariant, this reduces to $\iota_b^{Diff} = e^{-\hat{C}^*} \circ \iota_c^{Diff} \circ e^{\hat{C}_S^*}$. Composing (35) and (36) with the isomorphism $\underline{\text{Cyl}}_{\text{diff}}^* \leftrightarrow \text{Cyl}_{\text{diff}}^*$ defined in section 3 then provides us with the formal expression for the ‘c’ embedding into $\text{Cyl}_{\text{diff}}^*$, and for the ‘b’ embeddings into $\text{Cyl}_{\text{diff}}^*$. The use of $\underline{\text{Diff}}$ instead of Diff not only has allowed us to write these expressions, but the fact that $\underline{\text{Diff}}$ is so much smaller than Diff makes it more likely that they can be regularized.

6 Discussion

The kinematics of LQG are usually formulated in terms of the piecewise analytic category. We have shown that the piecewise analytic category is not essential, and can be replaced with something as simple as the piecewise linear category, giving rise to what we have called *piecewise linear LQG* (pLQG). We have shown that piecewise linear LQG is fully equivalent to standard LQG at the diffeomorphism invariant level, both in terms of Hilbert space structure and dynamics, as long as one makes a natural choice of generalized diffeomorphism group such as advocated in [13].

Furthermore, we have seen that LQC is exactly embeddable into pLQG. This shows that the non-embeddability result of [6] is perhaps somewhat of a red herring: it appears relevant at the kinematical level, but this relevance seems to evaporate at the diffeomorphism invariant level. For, pLQG circumvents the non-embeddability result of [6], and is yet fully equivalent to LQG at the diffeomorphism invariant level. This is what has now allowed us to at least write down formal expressions for embeddings of LQC into LQG at the diffeomorphism invariant level. These expressions were given in section 5.¹¹ Of course it still remains to regulate these expressions in some way.

¹¹As a side note, it may also be possible that there is another way to relate LQG to cosmology other than via the piecewise linear framework presented here. For, as pointed out by Koslowski [22], it appears that, given any analytic edge e , the holonomy along e as a function of the symmetric connection $A_S = r(c) = c \hat{A}_a^i$ can be decomposed into an almost periodic part [20] and a part vanishing as c approaches infinity. If true, it is not hard to see that this decomposition must be unique, as there are no almost periodic functions that vanish at infinity. This would then allow one to construct a projector $P_{\text{ap}} : r^*[\text{Cyl}] \rightarrow \text{Cyl}_S$ that projects out the almost periodic part. The projector could then be used to construct embeddings ι_c and ι_b of LQC directly into Cyl^* : $(\iota_c \psi | \Phi) := (\psi | P_{\text{ap}} r^* | \Phi)$, and then $\iota_b := e^{-\hat{C}^*} \circ \iota_c \circ e^{\hat{C}_S^*}$. These embeddings would again satisfy the physical intertwining criterion used in [5]. As the codomain of such embeddings would be directly Cyl^* , and Diff acts on Cyl^* , one would then be able to directly write down a formal expression for ‘c’ and ‘b’ embeddings into diffeomorphism invariant states, similar to that in section 5 of this paper. One could then check whether the resulting formal embedding is equivalent to the one given in this paper. Of course, the resulting embedding would also have to be regularized. For the present, this is just a future possibility.

We close with some remarks regarding the similarities of piecewise linear LQG to the framework underlying the construction of spinfoams. As argued, for example, in [9], the classical theory underlying spinfoams is a certain discrete theory based on piecewise flat geometries. Furthermore, as touched upon in appendix B of [9], in order for the discrete variables to fully describe the piecewise flat geometry, one is implicitly assuming a given linear structure on each patch. Thus, one is actually assuming a piecewise linear structure of spacetime. As seen in this paper, the use of piecewise linear structures naturally leads to the use of simplicial complexes, and simplicial complexes are central in the classical discrete theory underlying spinfoams. Whether the relation between pLQG and spinfoams goes beyond these cursory remarks is not clear, and would be interesting to investigate.

Acknowledgements

The author thanks Tim Koslowski for stimulating exchanges, Andrew Ranicki for encouraging him to study simplicial complexes in more depth, and Thomas Thiemann for remarks on a prior draft. This work was supported in part by the Alexander von Humboldt foundation of Germany.

A Existence of triangulation compatible with a graph

We prove here a result that is needed in section 3 for proving the unitary isomorphism between the diffeomorphism invariant Hilbert spaces of pLQG and LQG. We place it in an appendix because it requires a number of new definitions that are not needed elsewhere in the paper, and would distract from the logic of section 3.

For the purposes of this appendix, we remind the reader, from section 3, that a 1-complex X is said to be *compatible* with a piecewise linear graph γ if $|X|$ is equal to the image of γ . In the following, we will also need the notion of a *cell* and a *cell complex*, which we define here. To summarize in short, a *cell* is a compact convex polyhedron (see [23] for a definition in terms of more rudimentary notions). Given a cell C , and a plane P such that $C \setminus P$ is connected, we call $A := P \cap C$ a *face* of C , and we write $A < C$. The vertices, edges, faces of C in the usual sense, as well as C itself, are all faces of C . A *cell complex* is defined in a manner analogous to a simplicial complex:

Definition (cell complex). A *cell complex* K is a finite collection of cells satisfying (i) if $C \in K$ and B is a face of C , then $B \in K$ and (ii) If $B, C \in K$, then $B \cap C$ is a face of B and C .

Given a cell complex K , we define $|K| := \cup_{A \in K} A$ as the polyhedron underlying K . Given two points p, q , let pq denote the line segment between them. Then given a cell A and a point p not in the plane determined by A , one defines the *cone* with vertex p and base A , denoted pA , by $pA := \cup_{q \in A} (pq)$. Given two cell complexes K and L , K is said to be a *subdivision* of L if $|K| = |L|$ and every cell in K is contained in a cell of L . A subdivision K of L is said to be obtained by *starring* at a point a if K is obtained from L by replacing each cell $C \in L$ with $a \in C$ by the collection of cells $\{aF | F < C, a \notin F\}$ (see p.15 of [23]).

With these preliminaries out of the way, we come to the theorem.

Theorem 12. *Given any piecewise linear graph γ , there exists a triangulation K of $\mathbb{R}^3$ containing a 1-dimensional subcomplex K_1 compatible with γ .*

Proof. Let X be the minimal 1-complex compatible with γ : that is, break up each edge of γ into its straight parts, and then define X to contain all of these straight parts and their end points. Let N denote any rectangular prism sufficiently large so that it contains all of X , without X intersecting the boundary of N .

For each 1-simplex e in X , let e_1, e_2 denote the end points. Construct a cell complex N_{e_1} by starring N at the point e_1 . Then construct N_e by starring N_{e_1} at e_2 . Because every cell of N_{e_1} containing e_2 possesses e_1 as a vertex, the starring procedure gaurantees $e_2 e_1 = e$ will belong to N_e . Take the repeated *intersection* of the cell complexes N_e ,

$$Q := \{\cap_{e \in X} A_e \mid \{A_e \in N_e\}_{e \in X}\}. \quad (37)$$

Noting example 2.8(5) of [23], this is again a cell complex. Furthermore, Q is a subdivision of each cell complex A_e . It therefore contains a subdivision $\tilde{e}$ of each e . Taking the union of these subdivisions $\tilde{e}$ provides a 1-complex K_1 that is a subcomplex of Q , and that is compatible with γ . Next, from proposition 2.9 of [23], Q can be subdivided further to obtain a simplicial complex H , without adding any vertices, so that K_1 is again a subcomplex of H . Now, $|H| = N$ is a rectangular prism. Choose a vertex v of N , and let P_1, P_2, P_3 denote the three planes passing through v that contain two dimensional faces of N . By reflecting H repeatedly across these three planes, we obtain seven further copies of H that match on their common boundaries among themselves and with H . The union of H with these copies therefore defines a simplicial complex J . J is furthermore such that if we introduce an infinite number of copies of it, tiling all of $\mathbb{R}^3$, these copies will match on their common boundaries. If we let K denote the union of J with this infinite number of copies of J , then K is a simplicial complex. K triangulates all of $\mathbb{R}^3$, and contains the one dimensional subcomplex K_1 , which is compatible with γ . ■

References

- [1] Ashtekar A and Lewandowski J 2004 Background Independent Quantum Gravity: A Status Report *Class. Quant. Grav.* **21** R53.
- [2] Rovelli C 2004 *Quantum Gravity* (Cambridge: Cambridge UP).
 Smolin L 2004 An invitation to loop quantum gravity, in *Cincinnati 2003, Quantum theory and symmetries* pp 655-682 (*Preprint: hep-th/0408048*).
 Thiemann T 2003 Lectures on loop quantum gravity *Lect. Notes. Phys.* **631** 41-135 (*Preprint: gr-qc/0210094*).
 Thiemann T 2007 *Modern Canonical Quantum General Relativity* (Cambridge: Cambridge UP).
 Ashtekar A 2007 An introduction to loop quantum gravity through cosmology *Nouvo Cim.* **122B** 135-155.
- [3] Rovelli C and Smolin L 1988 Knot theory and quantum gravity *Phys. Rev. Lett.* **61** 1155-1158
 Rovelli C and Smolin L 1990 Loop representation for quantum general relativity *Nucl. Phys.* B331 80-152
 Ashtekar A and Isham C J 1992 Representations of the holonomy algebras of gravity and non-Abelian gauge theories *Class. Quantum Grav.* **9** 1433-1468
 Ashtekar A and Lewandowski J 1993 Representation theory of analytic holonomy C* algebras, in *Knots and Quantum Gravity*, ed. J Baez (Oxford: Oxford UP)
 Rovelli C and Smolin L 1995 Spin networks and quantum gravity *Phys. Rev. D* **52** 5743-5759
- [4] Engle J 2006 Quantum field theory and its symmetry reduction *Class. Quantum Grav.* **23** 2861-2894.
- [5] Engle J 2007 Relating loop quantum cosmology to loop quantum gravity: Symmetric sectors and embeddings *Class. Quant. Grav.* **24** 5777-5802.
- [6] Brunnemann J and Fleischhack C 2007 On the configuration spaces of homogeneous loop quantum cosmology and loop quantum gravity *arXiv:0709.1621*.

- [7] Zapata J 1997 A combinatorial approach to diffeomorphism invariant quantum gauge theories *J. Math. Phys.* **38** 5663-5681.
- [8] Reisenberger M P and Rovelli C 1997 ‘Sum over surfaces’ form of loop quantum gravity *Phys. Rev. D* **56** 3490-3508.
 Reisenberger M P 1997 A lattice worldsheet sum for 4-d euclidean GR, [arXiv: gr-qc/9711052](#).
 Baez J 1998 Spin foam models *Class. Quant. Grav.* **15** 1827-1858.
 Oriti D 2001 Spacetime geometry from algebra: spin foam models for non-perturbative quantum gravity *Rep. Prog. Phys.* **64** 1703-1757.
 Perez A 2003 Spin-foam models for quantum gravity *Class. Quant. Grav.* **20** R43-R104.
 Freidel L 2005 Group Field Theory: An overview *Int. J. Theor. Phys.* **44** 1769-1783.
- [9] Engle J, Pereira R and Rovelli C 2008 Flipped spinfoam vertex and loop gravity *Nucl. Phys.* **B789**.
- [10] Rovelli C and Smolin L 1995 Discreteness of area and volume in quantum gravity *Nucl. Phys.* **B442** 593-622.
 Ashtekar A and Lewandowski J 1996 Quantum theory of geometry I: Area operators *Class. Quantum Grav.* **14** A55-A82
 Ashtekar A and Lewandowski J 1998 Quantum theory of geometry II: Volume operators *Adv. Theor. Math. Phys.* **1**
- [11] Ashtekar A, Lewandowski J, Marolf D, Mourão J, and Thiemann T 1995 Quantization of diffeomorphism invariant theories of connections with local degrees of freedom *J. Math. Phys.* **36** 6456-6493.
- [12] Munkres J 1984 *Elements of Algebraic Topology* (Perseus Publishing: Cambridge, Massachusetts).
- [13] Koslowski T 2006 Physical Diffeomorphisms in Loop Quantum Gravity, [arXiv:gr-qc/0610017](#).
- [14] Fleischhack C 2006 Representations of the Weyl algebra in quantum geometry, [arXiv:math-ph/0407006](#).
- [15] Hardt R M 1975 Stratification of real analytic mappings and images, *Invent. Math.* **28** 193-208.
- [16] Fairbairn W and Rovelli C 2004 Separable Hilbert space in loop quantum gravity *J. Math. Phys.* **45** 2802-2814.
- [17] Brown E 1969 The hauptvermutung for 3-complexes *Trans. Amer. Math. Soc.* **144** 173-196.
- [18] Thiemann T 2006 Quantum spin dynamics VIII. The Master Constraint *Class. Quantum Grav.* **23** 2249-2266.
- [19] Thiemann T 1998 Quantum spin dynamics (QSD) *Class. Quantum Grav.* **15** 839-873.
- [20] Ashtekar A, Bojowald M and Lewandowski J 2003 Mathematical structure of loop quantum cosmology *Adv. Theor. Math. Phys.* **7** 233-268.
- [21] Thiemann T 2006 Complexifier coherent states for quantum general relativity *Class. Quantum Grav.* **23** 2063-2118.
- [22] Koslowski T, private communication.
- [23] Rourke C P and Sanderson B J 1982 *Introduction to Linear Topology* (Berlin : Springer-Verlag).